

OCCURRENCE OF "INDUCED" ANTIGENS IN SALMONELLAE
ISOLATED FROM MAN

P. R. EDWARDS AND MARY A. FIFE

*Communicable Disease Center, Public Health Service, U. S. Department of Health, Education, and Welfare,
Atlanta, Georgia*

Received for publication February 13, 1957

It is well known that motile cultures of salmonellae when inoculated into semisolid medium containing appropriate H antisera to prevent migration, may develop "induced" or "artificial" H antigens. Organisms which possess these changed antigens spread throughout the medium since the flagella no longer are influenced by the sera which immobilized the original cultures. Among the H antigens most prone to give rise to such changed forms is antigen b, which is changed to an antigen called z_{33} (Edwards and Moran, Proc. Soc. Exptl. Biol. Med., **61**, 242, 1946). This change is irrespective of the serotype in which the b antigen occurs. Thus, a monophasic form of *Salmonella minnesota* (21: b) placed in semisolid medium with anti-b serum will yield a 21: z_{33} form, whereas *Salmonella paratyphi* B (4,5,12: b-1,2) exposed to b and 1,2 sera is likely to yield 4,5,12: z_{33} — 1,2. Other H antigens yield characteristic "induced" forms when placed in contact with appropriate sera (Edwards and Bruner, J. Bacteriol., **38**, 63, 1939).

The natural occurrence of "induced" antigens has been noted only once, when Edwards and Moran (Proc. Soc. Exptl. Biol. Med., **61**, 242, 1946) recognized the 21: z_{33} form among cultures isolated from sewage by A. A. Hajna. Attention is called here to the appearance of these changed forms in man, including persons affected with gastroenteritis.

Within a period of 2 weeks in May 1950, 9 cultures were received from Miss Janie Morris and

Miss Alice Brim of the Georgia State Department of Health, to whom we are indebted for the following information: Four of the cultures were from persons in the same town. Three of the cultures were isolated from 3 adults who denied having any recent gastrointestinal symptoms. The fourth culture was isolated from the stool of a 12-year-old boy who had mild diarrhea.

The fifth person from whom the organisms were isolated was a 4-year-old boy with severe diarrhea who resided in a second town. The sixth isolation was from a person of unknown age and sex who was affected with diarrhea when admitted to a state tuberculosis sanatorium. The seventh and eighth cultures were isolated from the stools of 2 asymptomatic female workers in a school lunch room in a fourth town. The ninth culture was isolated in the course of a routine examination of the stools of a known typhoid carrier.

In agglutinin-adsorption tests the organisms were agglutinated with a 21: z_{33} serum, and the H and O agglutinins were removed. The serum had been derived from a monophasic culture of *S. minnesota* (21: b), which had been passed through b serum and thus changed to 21: z_{33} . The cultures were not flocculated by sera for the normal H antigens of *S. minnesota* (b and e,n,x).

The observations indicate the importance of recognizing the changes that may take place in H antigens of known *Salmonella* serotypes which otherwise might be designated as independent types. The "induced" form has not been recognized in the area since.

3-HYDROXY-2-NAPHTHOIC ACID AS AN INTERMEDIATE
IN BACTERIAL DISSIMILATION OF ANTHRACENE

MARTIN H. ROGOFF AND IRVING WENDER

Central Experiment Station, Bureau of Mines, U. S. Department of the Interior, Bruceton, Pennsylvania

Received for publication March 4, 1957

The authors have previously reported (Rogoff and Wender, J. Bacteriol., **73**, 264, 1957) the

isolation of 1-hydroxy-2-naphthoic acid as an intermediate in the breakdown of phenanthrene




Figure 1. Oxidation of aromatic compounds by anthracene-grown cells.




Figure 2. Pathways of polycyclic hydrocarbon oxidation.

by a bacterium resembling *Pseudomonas aeruginosa*. This product results from attack at an end ring as opposed to splitting of the reactive 9:10-bond as postulated by Tausson (Planta, **5**, 239, 1938). Studies on dissimilation of anthracene

were carried out to determine the mode of attack on a linearly condensed polycyclic system as opposed to the angularly condensed phenanthrene molecule.

The anthracene-decomposing organism, a gram negative motile rod, was isolated from soil by enrichment culture with naphthalene as a source of carbon. The organism closely resembles *Pseudomonas aeruginosa* and differs from the phenanthrene strain only in its inability to produce acid from glucose. Growth of the culture in serial transfers in liquid mineral-salts medium containing anthracene as a source of carbon, indicated the ability of the organism to utilize anthracene. After an incubation period of 2 to 3 weeks at room temperature on a shaker, aliquots of anthracene fermentations gave a faint green color with ferric chloride, indicating a phenolic intermediate. In several experiments, pairs of 4-L flasks containing 800 ml of mineral-salts medium and 0.5 per cent anthracene were inoculated with 100 ml of a broth culture, followed by incubation for 10 days on a rotary shaker (80-90 rpm). The cultures were removed from the shaker and allowed to stand at room temperature until aliquots gave a green color with ferric chloride, usually after an additional 2-3 weeks. The cultures were then neutralized with 5.0 per cent NaHCO_3 and concentrated under reduced pressure to one-tenth volume. After acidification with 1:1 HCl , the residues were continuously extracted with ether for 48 hr. The ether solutions were extracted with 5.0 per cent aqueous NaHCO_3 . The aqueous layer was separated, acidified, and centrifuged. The solid material obtained was recrystallized twice from hot water, yielding small amounts of yellow crystals. This material was identified as 3-hydroxy-2-naphthoic acid by comparison of its infrared and ultraviolet spectra with those of a known sample of the acid.

Manometric experiments with anthracene-grown cells demonstrated that the organism possessed the enzymes necessary to oxidize salicylic acid and catechol (figure 1). The isolation of 3-hydroxy-2-naphthoic acid from anthracene breakdown, and of 1-hydroxy-2-naphthoic acid from phenanthrene breakdown indicate end ring attack as a mode of oxidation of polycyclic systems by members of the genus *Pseudomonas*. The pathways diagrammed in figure 2 are tentatively offered for phenanthrene and anthracene dissimilation.